

MULTICOLUMN SCS FOR PREDOMINANT BACK PAIN IN FBSS PATIENTS



RIGOARD, PHILIPPE, ET AL. PAIN (2019).

218 PATIENTS

Randomized to SCS with optimal medical management (OMM) vs. OMM alone



Primary outcome: proportion of patients with >50% reduction in LBP at 6 months




NORTH AMERICAN AND EUROPEAN SITES



28

Prospective, open-label, parallel-group, randomized trial

Mean Low Back Pain Score by Treatment Group



* = statistical significance

SCS FOR PREDOMINANT LOW BACK PAIN



Adding multicolumn SCS to OMM improved pain relief, health-related quality of life and function in FBSS patients with predominant LBP, and sustained for 12 months

MULTICOLUMN SPINAL CORD STIMULATION FOR PREDOMINANT BACK PAIN IN FAILED BACK SURGERY SYNDROME PATIENTS: A MULTICENTER RANDOMIZED CONTROLLED TRIAL

Philippe Rigoard, Surajit Basu, Mehul Desai, Rod Taylor, Lieven Annemans, Ye Tan, Mary Jo Johnson, Carine Van den Abeele, PROMISE Study Group, Richard North

DOI: 10.1097/j.pain.0000000000001510

email us at info@neuromodulation.org